arXiv:nlin/0301011v1 [nlin.SI] 10 Jan 2003

General solution for Hamiltonians with extended cubic and quartic potentials *

C. Verhoeven, M. Musette and R. Conte

10 January 2003

Keywords. Hénon-Heiles, Hamiltonian systems, separation of variables, nonlinear equations, hyperelliptic functions, soliton equations.

Abstract

We integrate with hyperelliptic functions a two-particle Hamiltonian with quartic potential and additionnal linear and nonpolynomial terms in the Liouville integrable cases 1 : 6 : 1 and 1 : 6 : 8.

1 Introduction

The generalized Hénon-Heiles Hamiltonian

$$H = \frac{1}{2}(P_X^2 + P_Y^2 + c_1 X^2 + c_2 Y^2) + aXY^2 - \frac{b}{3}X^3 + \mu Y^{-2}, \quad \mu \text{ arbitrary} \qquad (1)$$

is known to be Liouville integrable [1, 2, 3] for three sets of values of $(b/a, c_1, c_2)$ and to be related [4] to the stationary reduction of the following fifth order soliton equations: KdV$_5$, Sawada-Kotera (SK) and Kaup-Kupershmidt (KK).
The canonical transformation [5] between the SK and KK cases for $\mu \neq 0$ allowed us [6] to define the separating variables of the Hamilton-Jacobi equations and to derive the general solution of the equations of motion with hyperelliptic functions.
The two-particle Hamiltonian with quartic potential

$$H = \frac{1}{2}(P_X^2 + P_Y^2) - \frac{1}{2}(aX^2 + bY^2) + CX^4 + BX^2Y^2 + AY^4, \qquad (2)$$

is known to be Liouville integrable in four cases [7, 8, 9]:

1. $A : B : C = 1 : 2 : 1, a, b$ arbitrary,
2. $A : B : C = 1 : 6 : 1, a = b$,
3. $A : B : C = 1 : 6 : 8, a = 4b$,

---

[*] *Theoretical and Mathematical Physics*, to appear. S2002/094. nlin.SI/0301011

4. $A : B : C = 1 : 12 : 16, a = 4b$.

The extension of (2) to include some linear and non polynomial terms which preserve the Liouville integrability [10, 9] and confirm their connection with some integrable soliton equations has been considered by various authors [11, 12].
The case 1 with extra inverse square terms is equivalent to the travelling wave reduction of the Manakov system [13] of two coupled NLS equations. It corresponds to the particular case for two particles of the extended Garnier system [14] which has been integrated by Wojciekowski [15]. The case 4 can be associated [5] with a coupled KdV system possessing a fifth order Lax pair [16].
The authors of [17, 5] have showed how cases 2 and 3 with some extra linear and inverse square terms are equivalent to the stationary reduction $\xi = x - ct$ of the Hirota-Satsuma system [18]:

$$r_t = \frac{1}{2} r_{xxx} + 3rr_x - 6ss_x, \quad s_t = -s_{xxx} - 3rs_x \tag{3}$$

and an other coupled KdV system

$$\left\{ \begin{array}{ll} f_t = & -\frac{1}{2}(2f_{xxx} + 3ff_{xx} + 3f_x^2 - 3f^2 f_x + 6fg_x + 6gf_x), \\ g_t = & \frac{1}{4}(2g_{xxx} + 12gg_x + 6fg_{xx} + 12gf_{xx} + 18f_x g_x - 6f^2 g_x \\ & +3f_{xxxx} + 3ff_{xxx} + 18f_x f_{xx} - 6f^2 f_{xx} - 6ff_x^2), \end{array} \right. \tag{4}$$

In the present paper, using the canonical transformation [5] between these two extended cases, we show how the method we followed for the SK and KK cases of (1) to define the separating variables of the Hamilton-Jacobi equation can be applied here. Then the equations of motion associated with cases 2 and 3 of (2) with extra linear and inverse square terms are also integrated with hyperelliptic functions.
In section 2, we consider the SK and KK integrable cases of the Hénon-Heiles Hamiltonian (1) for $\mu \neq 0$ to explain the method we applied in [6] for integrating their equations of motion. In section 3 we consider the Hamiltonian (2) with extended quartic potential in case 2 and 3 and recall the canonical transformation [5], which relates the case $1 : 6 : 1$ of (2) (with extra terms of the form $\alpha X^{-2} + \beta Y^{-2}$) and the case $1 : 6 : 8$ (with extra terms like $\gamma X + \delta Y^{-2}$).
To the best of our knowledge, the results of sections 4 and 5 are new. In section 4, we explicitly integrate the equations of motion of the case $1 : 6 : 8$ for $\delta \neq 0, \gamma = 0$. In section 5 we integrate the $1 : 6 : 1$ case with $\alpha = \beta \neq 0$ and we use the canonical transformation to transport those solutions to the $1 : 6 : 8$ case for $\delta = 0, \gamma \neq 0$.

2 Integration of the cubic Hénon-Heiles Hamiltonian

Let us consider the SK and KK Hénon-Heiles Hamiltonians with their second integral, denoted as K_2^2 and k_2^2, and their equations of motion [10, 9]:

$$\text{SK:} \qquad \frac{b}{a} = -1, \quad , c_1 = c_2, \quad a = \frac{1}{2}, \quad U = X + c_2, \quad V = Y, \quad c = c_1 c_2 \tag{5}$$

$$H \equiv K_1 = \frac{1}{2}(P_U^2 + P_V^2) + \frac{1}{2}UV^2 + \frac{1}{6}U^3 - \frac{c}{2}U + \frac{\mu}{8V^2}, \tag{6}$$

$$K_2^2 = K_{2,0}^2 + \frac{2}{3}\mu U + \mu\frac{P_U^2}{V^2}, \tag{7}$$

$$K_{2,0} = -2P_U P_V - U^2 V - \frac{V^3}{3} + cV, \tag{8}$$

$$U'' = -\frac{1}{2}(V^2 + U^2) - \frac{c}{2}, \qquad V'' = -UV + \frac{\mu}{4V^3}, \tag{9}$$

KK: $$\frac{b}{a} = -16, \quad c_1 = 16c_2, \quad a = \frac{1}{4}, \quad c = c_1 c_2, \quad u = X + 2c_2, \quad V = Y \tag{10}$$

$$H \equiv k_1 = \frac{1}{2}(p_u^2 + p_v^2) + \frac{1}{4}uv^2 + \frac{4}{3}u^3 - cu + \frac{1}{2}\frac{\mu}{v^2}, \tag{11}$$

$$k_2^2 = k_{2,0}^2 + \frac{\mu}{3}u + 2\mu\frac{p_v^2}{v^2} + \frac{\mu^2}{v^4}, \tag{12}$$

$$k_{2,0}^2 = p_v^4 - \frac{1}{72}v^6 - \frac{1}{12}u^2v^4 + up_v^2v^2 - \frac{1}{3}p_u p_v v^3 + \frac{c}{12}v^4, \tag{13}$$

$$u'' = -\frac{1}{4}v^2 - 4u^2 + c, \qquad v'' = -\frac{1}{2}uv + \frac{\mu}{v^3}, \tag{14}$$

which are equivalent to the stationary reduction $\xi = x - ct$ of the two integrable PDE's:

$$\text{SK:} \;\; U_t + (U_{xxxx} + 5UU_{xx} + \frac{5}{3}U^3)_x = 0, \tag{15}$$

$$\text{KK:} \;\; u_t + (u_{xxxx} + 10uu_{xx} + \frac{20}{3}u^3 + 30u_x^2)_x = 0. \tag{16}$$

Both equations are connected to another fifth order integrable PDE [19]:

$$w_t + (w_{4x} - 5w_x w_{xx} - 5w^2 w_{xx} - 5ww_x^2 + w^5)_x = 0. \tag{17}$$

by the Miura transformation

$$U = w_x - w^2, \qquad u = -w_x - \frac{1}{2}w^2. \tag{18}$$

Solving the stationary reduction of (17) for w in function of V, V' and v, v' and defining the following expressions:

$$\begin{aligned} \lambda^2 &= -\mu, \;\; \Gamma = 6(VK_{2,0} + \lambda P_U), \\ \Omega &= 48(3v^4k_{2,0}^2 + 6\lambda uv^5p_v + 12\lambda p_v^3v^3 - \lambda v^6 p_u \\ &\quad +3\lambda^2uv^4 + 18\lambda^2v^2p_v^2 + 12\lambda^3vp_v + 3\lambda^4), \end{aligned} \tag{19}$$

one obtains the following canonical transformation [5, 20]:

$$u = -\frac{3}{2}\left(-\frac{P_V}{V} + \frac{\lambda}{2V^2}\right)^2 - U, \quad v = \frac{\sqrt{\Gamma}}{V}, \tag{20}$$

$$\begin{aligned} p_u &= \frac{1}{V^3}(3P_V^3 + 3UV^2P_V - P_UV^3) \\ &\quad -\frac{3\lambda}{2V^6}\left(UV^4 + 3V^2P_V^2 - \frac{3}{2}\lambda V P_V + \frac{\lambda^2}{4}\right), \\ p_v &= \frac{1}{4V^2}\left(-2P_V + \frac{\lambda}{V}\right)\sqrt{\Gamma} - \lambda\frac{V}{\sqrt{\Gamma}}, \end{aligned} \tag{21}$$

$$U = -6\left(\frac{p_v}{v} + \frac{\lambda}{v^2}\right)^2 - u, \quad V = \frac{\sqrt{\Omega}}{2v^4}, \tag{22}$$

$$\begin{aligned} P_U &= \frac{1}{v^3}(12p_v^3 + 6uv^2p_v - v^3p_u) \\ &\quad +\frac{3\lambda}{v^6}(2uv^4 + 12v^2p_v^2 + 12\lambda v p_v + 4\lambda^2), \end{aligned} \tag{23}$$

$$P_V = -\frac{1}{v^5}\left(p_v + \frac{\lambda}{v}\right)\sqrt{\Omega} + \lambda\frac{v^4}{\sqrt{\Omega}}. \tag{24}$$

For $\mu = 0$, we introduce the expressions (22)–(23) for U, V, P_U, P_V in the variables which separate the SK Hamiltonian

$$\begin{aligned} Q_1 = U + V, \quad P_1 = \tfrac{1}{2}(P_U + P_V), \\ Q_2 = U - V, \quad P_2 = \tfrac{1}{2}(P_U - P_V). \end{aligned} \tag{25}$$

This defines the change of variables [21]

$$\begin{aligned} q_1 &= -6\frac{p_v^2 - k_{2,0}}{v^2} - u, \qquad p_1 = \frac{1}{2v^3}(12p_v^3 + 6uv^2p_v - v^3p_u - 12p_vk_{2,0}), \\ q_2 &= -6\frac{p_v^2 + k_{2,0}}{v^2} - u, \qquad p_2 = \frac{1}{2v^3}(12p_v^3 + 6uv^2p_v - v^3p_u + 12p_vk_{2,0}), \end{aligned} \tag{26}$$

that we apply on the KK Hamiltonian (11), taking account that for $\mu \neq 0$, $k_{2,0}$ is no more a constant of motion. Therefore, one has

$$H \equiv k_1 = p_1^2 + p_2^2 + \frac{1}{12}(q_1^3 + q_2^3) - \frac{c}{4}(q_1 + q_2) + \frac{\mu}{24}\frac{q_1 - q_2}{k_{2,0}}, \tag{27}$$

$$k_{2,0} = 2(p_2^2 - p_1^2) + \frac{1}{6}(q_2^3 - q_1^3) - \frac{c}{2}(q_2 - q_1), \tag{28}$$

$$q_1' = 2p_1 + \frac{\mu}{6}\frac{(q_1 - q_2)p_1}{k_{2,0}^2}, \tag{29}$$

$$q_2' = 2p_2 - \frac{\mu}{6}\frac{(q_1 - q_2)p_2}{k_{2,0}^2}. \tag{30}$$

In this new setting of coordinates, it is important to note that, introducing the expression

$$f(q_i, p_i) = 2p_i^2 + \frac{1}{6}q_i^3 - \frac{c}{2}q_i \text{ for } i = 1, 2, \tag{31}$$

in (27) and (28), the Hamilton-Jacobi equation is separated for μ arbitrary.
In the variables (26), the second invariant k_2^2 can be written in two equivalent ways:

$$k_2^2 = -\frac{\mu}{3} q_1 + \left(k_{2,0} + \frac{\mu}{12} \frac{q_1 - q_2}{k_{2,0}} \right)^2, \tag{32}$$

$$\text{or} \qquad k_2^2 = -\frac{\mu}{3} q_2 + \left(k_{2,0} - \frac{\mu}{12} \frac{q_1 - q_2}{k_{2,0}} \right)^2, \tag{33}$$

which, after the elimination of $\mu(q_1 - q_2)/k_{2,0}$ between (27)–(32) and (27)–(33), become

$$k_2^2 = -\frac{\mu}{3} q_1 + \left(-4p_1^2 - \frac{q_1^3}{3} + c q_1 + 2k_1 \right)^2, \tag{34}$$

$$\text{or} \qquad k_2^2 = -\frac{\mu}{3} q_2 + \left(4p_2^2 + \frac{q_2^3}{3} - c q_2 - 2k_1 \right)^2 \cdot \tag{35}$$

Then we can eliminate p_1 between (34) and (29), and p_2 between (35) and (30) to obtain

$$q_1' = \sqrt{2k_1 - \frac{q_1^3}{3} + c q_1 - \sqrt{k_2^2 + \frac{\mu}{3} q_1}} \left(1 + \frac{\mu}{3} \frac{q_1 - q_2}{\left(\sqrt{k_2^2 + \frac{\mu}{3} q_2} + \sqrt{k_2^2 + \frac{\mu}{3} q_1} \right)^2} \right), \tag{36}$$

$$q_2' = \sqrt{2k_1 - \frac{q_2^3}{3} + c q_2 + \sqrt{k_2^2 + \frac{\mu}{3} q_2}} \left(1 - \frac{\mu}{3} \frac{q_1 - q_2}{\left(\sqrt{k_2^2 + \frac{\mu}{3} q_2} + \sqrt{k_2^2 + \frac{\mu}{3} q_1} \right)^2} \right) \cdot \tag{37}$$

For $\mu \neq 0$, setting:

$$s_1 = \sqrt{3\frac{k_2^2}{\mu} + q_1}, \qquad s_2 = -\sqrt{3\frac{k_2^2}{\mu} + q_2}, \tag{38}$$

and defining

$$P(s) = 2k_1 - \frac{1}{3} \left(s^2 - 3\frac{k_2^2}{\mu} \right)^3 + c \left(s^2 - 3\frac{k_2^2}{\mu} \right) - \sqrt{\frac{\mu}{3}} s, \tag{39}$$

the equations (36)–(37) become:

$$s_1' = \frac{\sqrt{P(s_1)}}{s_1 - s_2}, \; s_2' = -\frac{\sqrt{P(s_2)}}{s_1 - s_2} \cdot \tag{40}$$

They can be solved by inverting the hyperelliptic integrals

$$\int_\infty^{s_1} \frac{\mathrm{d}s}{\sqrt{P(s)}} + \int_\infty^{s_2} \frac{\mathrm{d}s}{\sqrt{P(s)}} = k_3, \tag{41}$$

$$\int_\infty^{s_1} \frac{s\mathrm{d}s}{\sqrt{P(s)}} + \int_\infty^{s_2} \frac{s\mathrm{d}s}{\sqrt{P(s)}} = \xi + k_4. \tag{42}$$

and the general solution for the KK system is

$$u = -\frac{1}{2}(s_1^2 + s_2^2) + \frac{3}{\mu}k_2^2 - \frac{3}{2}\left(\frac{s_1' + s_2'}{s_1 + s_2}\right)^2, \qquad v^2 = \frac{2\sqrt{3\mu}}{s_1 + s_2}\cdot \tag{43}$$

For $\mu = 0$ from (36)–(37) and (26), one easily recovers the known solution [21] expressed with Weierstrass elliptic functions:

$$u = 6(\wp_1 + \wp_2) - \frac{3}{2}\left(\frac{\wp_1' - \wp_2'}{\wp_1 - \wp_2}\right)^2, \qquad v^2 = \frac{k_{2,0}}{\wp_2 - \wp_1}, \tag{44}$$

where $\wp_i$ satisfies the equation

$$\begin{aligned} &\wp_i'^2 = 4\wp_i^3 - g_2\wp_i - g_3^{(i)}, \quad i = 1, 2 && (45)\\ &g_2 = \frac{c}{12}, \qquad g_3^{(1)} = -\frac{1}{144}(2k_{1,0} - k_{2,0}), \quad g_3^{(2)} = -\frac{1}{144}(2k_{1,0} + k_{2,0}). \end{aligned}$$

In the SK case, from (43) and the canonical transformation (22), the general solution writes

$$\begin{aligned} &U = \sqrt{-3}(s_1' + s_2') + s_1^2 + s_1 s_2 + s_2^2 - \tfrac{3}{\mu}K_2^2,\\ &V^2 = -2\sqrt{-3}(s_1 + s_2)(s_1 s_1' + s_2 s_2') + 2(s_1 + s_2)^2\left(s_1^2 + s_2^2 - \tfrac{9K_2^2}{2\mu}\right), \end{aligned} \tag{46}$$

which, in the particular case $\mu = 0$, is merely

$$U = -6\left(\wp_1 + \wp_2\right), \qquad V = -6\left(\wp_1 - \wp_2\right)\cdot \tag{47}$$

3 Canonical transformation between extended $1:6:1$ and $1:6:8$ Hamiltonians

Let us consider the quartic Hamiltonian (2) with extra linear and inverse quadratic terms in the two following cases [9]:

$$\begin{aligned} 1:6:1 \quad H \equiv K_1 = {} & \frac{1}{2}(P_U^2 + P_V^2) - \frac{1}{32}(U^4 + 6U^2V^2 + V^4)\\ & -\frac{c}{2}(U^2 + V^2) + \frac{1}{2}(\frac{\alpha}{U^2} + \frac{\beta}{V^2}), \end{aligned} \tag{48}$$

$$K_2^2 = K_{2,0}^2 - \beta U^2 - \alpha V^2 + 4(\alpha\frac{P_V^2}{U^2} + \beta\frac{P_U^2}{V^2}) + \frac{4\alpha\beta}{U^2V^2} - 4c(\alpha + \beta), \tag{49}$$

$$K_{2,0} = 2P_U P_V - \frac{1}{4}UV(U^2 + V^2 + 8c), \tag{50}$$

$$U' = P_U, \qquad V' = P_V, \tag{51}$$

$$U'' = \frac{1}{8}U^3 + \frac{3}{8}UV^2 + cU + \frac{\alpha}{U^3} \quad V'' = \frac{1}{8}V^3 + \frac{3}{8}U^2V + cV + \frac{\beta}{V^3}, \tag{52}$$

$$\begin{aligned} 1:6:8 \quad H \equiv k_1 &= \frac{1}{2}(p_u^2+p_v^2) - \frac{1}{16}(8u^4+6u^2v^2+v^4) \\ &\quad -\frac{c}{2}(4u^2+v^2) - \gamma u + \frac{\delta}{2v^2}, \end{aligned} \tag{53}$$

$$\begin{aligned} k_2^2 = {} & k_{2,0}^2 + \tfrac{\delta^2}{v^4} - \gamma v^2 u^3 - \tfrac{\gamma}{2} u v^4 - \tfrac{\delta}{2} u^2 - 2\gamma^2 v^2 + 2\delta \tfrac{p_v^2}{v^2} - \tfrac{\delta}{4} v^2 \\ & +4\gamma(v p_u p_v - u p_v^2) - 4\gamma\delta \tfrac{u}{v^2} - 2\delta c - 4c\gamma v^2 u, \end{aligned} \tag{54}$$

$$\begin{aligned} k_{2,0}^2 = {} & (-p_v^2 + \tfrac{1}{8}v^4 + \tfrac{1}{4}u^2v^2 - uvp_v + \tfrac{1}{2}p_u v^2 + cv^2) \\ & \times(-p_v^2 + \tfrac{1}{8}v^4 + \tfrac{1}{4}u^2v^2 + uvp_v - \tfrac{1}{2}p_u v^2 + cv^2), \end{aligned} \tag{55}$$

$$u' = p_u, \qquad v' = p_v, \tag{56}$$

$$u'' = 2u^3 + \frac{3}{4}uv^2 + 4cu + \gamma, \quad v'' = \frac{1}{4}v^3 + \frac{3}{4}u^2 v + cv + \frac{\delta}{v^3}, \tag{57}$$

where K_1, K_2 and k_1, k_2 are the constants of motion of both systems.
The equations of motion are respectively equivalent to the stationary reduction $\xi = x - ct$ of the PDE's (3) and (4) with the correspondence:

$$r = -\frac{1}{4}(U^2+V^2+4c), \qquad s = \frac{1}{8}(U^2-V^2), \tag{58}$$

$$f = u, \qquad g = -\frac{1}{4}(v^2+2u^2+2p_u+4c)\cdot \tag{59}$$

Both PDE's (3) and (4) possess a fourth order Lax pair with scattering operators L and $\widetilde{L}$ which factorize as follows [5, 17],

$$\begin{aligned} 1:6:1 \quad L &= (\partial_x^2 + r + s)(\partial_x^2 + r - s) \\ &= (\partial_x - v_1)(\partial_x + v_1)(\partial_x + v_2)(\partial_x - v_2), \end{aligned} \tag{60}$$

$$\begin{aligned} 1:6:8 \quad \widetilde{L} &= (\partial_x^2 + f\partial_x + f_x + g)(\partial_x - f\partial_x + g) \\ &= (\partial_x + v_1)(\partial_x + v_2)(\partial_x - v_2)(\partial_x - v_1), \end{aligned} \tag{61}$$

and yield the Miura maps:

$$1:6:1 \quad r = \frac{1}{2}(v_{1,x} - v_{2,x} - v_1^2 - v_2^2), \quad s = \frac{1}{2}(v_{1,x} + v_{2,x} - v_1^2 + v_2^2), \tag{62}$$

$$1:6:8 \quad f = v_1 + v_2, \qquad g = v_1 v_2 - v_{1,x}, \tag{63}$$

while v_1 and v_2 are solutions of the system of PDE's

$$\begin{aligned} v_{1,t} &= \tfrac{1}{8}(-2v_{1,xx} - 6v_{2,xx} - 12v_2v_{2,x} - 12v_1v_{2,x} - 12v_1v_2^2 + 4v_1^3)_x, \\ v_{2,t} &= \tfrac{1}{8}(-2v_{2,xx} - 6v_{1,xx} + 12v_1v_{1,x} + 12v_2v_{1,x} - 12v_2v_1^2 + 4v_2^3)_x. \end{aligned} \tag{64}$$

From the relations (62)–(63), the transformations (58)–(59) and the equations of motion (51)–(52), (56)–(57), the stationary reduction of (64) can be solved for v_1 and v_2 in function of the canonical variables of the two extended systems:

$$1:6:1 \qquad v_1 = -\frac{P_U}{U} + \frac{\sqrt{-\alpha}}{U^2}, \qquad v_2 = \frac{P_V}{V} + \frac{\sqrt{-\beta}}{V^2}, \tag{65}$$

$$1:6:8 \qquad v_1 = \frac{u}{2} + \frac{p_v}{v} + \frac{\sqrt{-\delta}}{v^2}, \qquad v_2 = \frac{u}{2} - \frac{p_v}{v} - \frac{\sqrt{-\delta}}{v^2}, \tag{66}$$

such that, defining

$$\alpha = -\kappa_1^2, \quad \beta = -\kappa_2^2, \quad \delta = -(\kappa_2 - \kappa_1)^2, \quad \gamma = \frac{1}{2}(\kappa_1 + \kappa_2), \tag{67}$$

$$\Omega = -\frac{1}{4}(U^2 + V^2) + \frac{2}{UV}(P_U P_V + \kappa_2 \frac{P_U}{V} - \kappa_1 \frac{P_V}{U} - \frac{\kappa_1 \kappa_2}{UV}) - 2c, \tag{68}$$

$$\begin{aligned}\Gamma_\mp = -4\Big(&\mp 2p_u + \frac{1}{2}v^2 + u^2 - 4\frac{p_v^2}{v^2} \pm 4\frac{up_v}{v} + 8(\kappa_2 - \kappa_1)\frac{p_v}{v^3} \\ &\mp 4(\kappa_2 - \kappa_1)\frac{u}{v^2} - 4\frac{(\kappa_2 - \kappa_1)^2}{v^4} + 4c\Big),\end{aligned} \tag{69}$$

the canonical transformation between the 1 : 6 : 1 and 1 : 6 : 8 cases becomes [5]

$$u = -\frac{P_U}{U} + \frac{P_V}{V} + \frac{\kappa_1}{U^2} + \frac{\kappa_2}{V^2},$$

$$p_u = -\frac{1}{4}(V^2 - U^2) + \frac{P_U^2}{U^2} - \frac{P_V^2}{V^2} - 2\kappa_1 \frac{P_U}{U^3} - 2\kappa_2 \frac{P_V}{V^3} + \frac{\kappa_1^2}{U^4} - \frac{\kappa_2^2}{V^4}, \tag{70}$$

$$v = 2\sqrt{\Omega} \qquad p_v = \sqrt{\Omega}\left(-\frac{P_U}{U} - \frac{P_V}{V} + \frac{\kappa_1}{U^2} - \frac{\kappa_2}{V^2}\right) - \frac{\kappa_2 - \kappa_1}{2\sqrt{\Omega}}, \tag{71}$$

$$U = \frac{1}{2}\sqrt{\Gamma_-}, \qquad P_U = -\frac{1}{2}\sqrt{\Gamma_-}\left(\frac{u}{2} + \frac{p_v}{v} - \frac{\kappa_2 - \kappa_1}{v^2}\right) + 2\kappa_1\sqrt{\Gamma_-}, \tag{72}$$

$$V = \frac{1}{2}\sqrt{\Gamma_+}, \qquad P_V = \frac{1}{2}\sqrt{\Gamma_+}\left(\frac{u}{2} - \frac{p_v}{v} + \frac{\kappa_2 - \kappa_1}{v^2}\right) - 2\kappa_2\sqrt{\Gamma_+}. \tag{73}$$

4 General solution of the $1:6:8$ Hamiltonian for $\gamma = 0$

For $\alpha = \beta = 0$, one introduces the transformation (72)–(73) in the variables

$$\begin{aligned} Q_1 &= \tfrac{1}{2}(U+V)^2, \quad P_1 = \frac{1}{2}\frac{P_U + P_V}{U + V}, \\ Q_2 &= \tfrac{1}{2}(U-V)^2, \quad P_2 = \frac{1}{2}\frac{P_U - P_V}{U - V}, \end{aligned} \tag{74}$$

which separated the 1 : 6 : 1 Hamiltonian (48) in the polynomial case. This defines the following change of variables:

$$q_1 = 4\frac{k_{2,0} + p_v^2}{v^2} - \frac{v^2}{2} - u^2 - 4c, \tag{75}$$

$$p_1 = \frac{-8vp_v p_u + 8up_v^2 + 2v^2u^3 + u(v^4 + 8k_{2,0}) + 8cuv^2}{16v(vp_u - 2up_v)}, \tag{76}$$

$$q_2 = 4\frac{-k_{2,0} + p_v^2}{v^2} - \frac{v^2}{2} - u^2 - 4c, \tag{77}$$

$$p_2 = \frac{-8vp_v p_u + 8up_v^2 + 2v^2u^3 + u(v^4 - 8k_{2,0}) + 8cuv^2}{16v(vp_u - 2up_v)}, \tag{78}$$

that we apply on the Hamiltonian system (53) for $\kappa_1\kappa_2 \neq 0$, i.e. when $k_{2,0}$ is no more a constant of motion. This transformation restricted to $\kappa_2 = -\kappa_1$ ($\delta = -4\kappa_1^2$, $\gamma = 0$) defines the coordinates which separate [11, 12] the Hamilton-Jacobi equation. Indeed, one has

$$H \equiv k_1 = \frac{1}{16}\Big(32q_1p_1^2 - q_1^2 + 32q_2p_2^2 - q_2^2 - 8c(q_1+q_2)\Big) - \kappa_1^2\frac{(q_1-q_2)}{4k_{2,0}}, \tag{79}$$

with

$$k_{2,0} = \frac{1}{8}\Big(32q_1p_1^2 - q_1^2 - 32q_2p_2^2 + q_2^2 + 8c(q_2-q_1)\Big), \tag{80}$$

$$q_1' = 4q_1p_1 + 2\kappa_1^2\frac{(q_1-q_2)q_1p_1}{k_{2,0}^2}, \tag{81}$$

$$q_2' = 4q_2p_2 - 2\kappa_1^2\frac{(q_1-q_2)q_2p_2}{k_{2,0}^2}, \tag{82}$$

such that defining

$$f(q_i,p_i) \equiv 32q_ip_i^2 - q_i^2 - 8cq_i, \quad i = 1,2$$

we have the following separated Hamilton-Jacobi equation:

$$f(q_1,p_1)^2 - f(q_2,p_2)^2 - 32\kappa_1^2(q_1-q_2) = 16k_1\left(f(q_1,p_1) - f(q_2,p_2)\right), \quad p_i = \frac{\partial S}{\partial q_i}. \tag{83}$$

In analogy with the Kaup-Kupershmidt case, we can write the second invariant in two equivalent ways:

$$k_2^2 = -2\kappa_1^2q_1 + \left(k_{2,0} + \frac{\kappa_1^2(q_1-q_2)}{2k_{2,0}}\right)^2, \tag{84}$$

$$\text{or} \qquad k_2^2 = -2\kappa_1^2q_2 + \left(k_{2,0} - \frac{\kappa_1^2(q_1-q_2)}{2k_{2,0}}\right)^2, \tag{85}$$

which allow us to eliminate $\kappa_1^2(q_1-q_2)/k_{2,0}$ between (79)–(84) and (79)–(85). Next, we eliminate p_1 and p_2 between those two resulting expressions and the equations (81), (82) and obtain:

$$q_1' = q_1\sqrt{\frac{q_1}{2} + 4c + 4\frac{k_1}{q_1} + \frac{2}{q_1}\sqrt{k_2^2+2\kappa_1^2q_1}}\left(1 + \frac{2\kappa_1^2(q_1-q_2)}{\left(\sqrt{k_2^2+2\kappa_1^2q_1}+\sqrt{k_2^2+2\kappa_1^2q_2}\right)^2}\right), \tag{86}$$

$$q_2' = q_2\sqrt{\frac{q_2}{2} + 4c + 4\frac{k_1}{q_2} - \frac{2}{q_2}\sqrt{k_2^2+2\kappa_1^2q_2}}\left(1 - \frac{2\kappa_1^2(q_1-q_2)}{\left(\sqrt{k_2^2+2\kappa_1^2q_1}+\sqrt{k_2^2+2\kappa_1^2q_2}\right)^2}\right). \tag{87}$$

For $\kappa_1 = 0$ the general solution of equations (86)–(87) can be expressed in terms of Weierstrass elliptic functions:

$$q_1 + \frac{8}{3}c = 8\wp\Big(\xi - \xi_1, \frac{4}{3}c^2 - \frac{k_1}{2} - \frac{k_2}{4}, \frac{c}{12}(2k_1 + k_2 - \frac{32}{9}c^2)\Big) \equiv 8\wp_1, \tag{88}$$

$$q_2 + \frac{8}{3}c = 8\wp\Big(\xi - \xi_2, \frac{4}{3}c^2 - \frac{k_1}{2} + \frac{k_2}{4}, \frac{c}{12}(2k_1 - k_2 - \frac{32}{9}c^2)\Big) \equiv 8\wp_2. \tag{89}$$

For $\kappa_1 \neq 0$ setting in (86)–(87):

$$s_1 = \sqrt{\frac{1}{2\kappa_1^2}k_2^2 + q_1} \quad \text{and} \quad s_2 = -\sqrt{\frac{1}{2\kappa_1^2}k_2^2 + q_2}, \tag{90}$$

and defining

$$\widetilde{P}(s) \equiv \frac{1}{2}\left(s^2 - \frac{k_2^2}{2\kappa_1^2}\right)^3 + 4c\left(s^2 - \frac{k_2^2}{2\kappa_1^2}\right)^2 + (4k_1 + 2\sqrt{2}\kappa_1 s)\left(s^2 - \frac{k_2^2}{2\kappa_1^2}\right), \tag{91}$$

we obtain the equations,

$$s_1' = \frac{\sqrt{\widetilde{P}(s_1)}}{s_1 - s_2},\, s_2' = -\frac{\sqrt{\widetilde{P}(s_2)}}{s_1 - s_2}, \tag{92}$$

which are solved with the inversion of the hyperelliptic integrals

$$\int_\infty^{s_1} \frac{\mathrm{d}s}{\sqrt{\widetilde{P}(s)}} + \int_\infty^{s_2} \frac{\mathrm{d}s}{\sqrt{\widetilde{P}(s)}} = k_3, \tag{93}$$

$$\int_\infty^{s_1} \frac{s\mathrm{d}s}{\sqrt{\widetilde{P}(s)}} + \int_\infty^{s_2} \frac{s\mathrm{d}s}{\sqrt{\widetilde{P}(s)}} = \xi + k_4\cdot \tag{94}$$

Therefore, for δ arbitrary, $\gamma = 0$, the general solution of the $1:6:8$ Hamiltonian is defined with symmetric combinations of s_1, s_2

$$u^2 = -\frac{1}{2}(s_1^2 + s_2^2) + \frac{k_2^2}{2\kappa_1^2} + \left(\frac{s_1' + s_2'}{s_1 + s_2}\right)^2 - \frac{2\sqrt{2}\kappa_1}{s_1 + s_2} - 4c, \quad v^2 = \frac{4\sqrt{2}\kappa_1}{s_1 + s_2}, \tag{95}$$

and is a single-valued function of the complex variable ξ. For $\kappa_1 = 0$ it degenerates into

$$u^2 = -4(\wp_1 + \wp_2) + \left(\frac{\wp_1' - \wp_2'}{\wp_1 - \wp_2}\right)^2 + \frac{k_{2,0}}{2(\wp_2 - \wp_1)} - \frac{4}{3}c, \qquad v^2 = \frac{k_{2,0}}{\wp_1 - \wp_2}\cdot \tag{96}$$

5 General solution of the extended $1:6:1$ Hamiltonian

Now, we again use the canonical transformation (72)–(73) to find the general solution of the $1:6:1$ Hamiltonian.
For $\kappa_1 = \kappa_2 = 0$, we introduce the expressions (96) in (72)–(73) and obtain the solution of the equations of motion

$$U^2 + V^2 = 8(\wp_1 + \wp_2 - \frac{2}{3}c), \qquad UV = 4(\wp_1 - \wp_2)\cdot \tag{97}$$

Next, for $\alpha\beta \neq 0$, with the restriction $\alpha = \beta$, we obtain the expressions

$$U^2 + V^2 = -2\sqrt{2}(s_1' + s_2') + 2(s_1^2 + s_2^2 + s_1 s_2) - \frac{K_2^2}{\kappa_1^2}, \tag{98}$$

$$\begin{aligned} U^2 V^2 &= -2\sqrt{2}(s_1 + s_2)(s_1 s_1' + s_2 s_2' - 2\kappa_1) \\ &\quad +2(s_1 + s_2)^2 \left(s_1^2 + s_2^2 - \frac{3}{4}\frac{K_2^2}{\kappa_1^2} + 4c \right), \end{aligned} \tag{99}$$

which depend on symmetric combinations of s_1, s_2 and therefore are single-valued functions of ξ.
Since the Hamiltonian (48) for $\alpha = \beta$ is an even function of κ_1, we apply on (98)–(99) the canonical transformation (70)–(71) for $\kappa_1 = \kappa_2$, and obtain the general solution of the 1 : 6 : 8 Hamiltonian in the extended case $\delta = 0$.

6 Conclusion

Romeiras [22] described a procedure which relates the extended cubic potential $\widetilde{V}_3(u, v) = V_3(u, v) + \mu v^{-2}/2$ in the KK case with the extended quartic potential $\widetilde{V}_4(u, v) = V_4(u, v) + \delta v^{-2}/2$ in the 1 : 6 : 8 case. This explains why the combination $u^2 + v^2/2 + 4c$ of the solutions (95) is identical to the solution (43). The question remains open to extrapolate the present results to the case $\alpha \neq \beta$, i.e. $\gamma\delta \neq 0$.

Acknowledgments

CV, a research assistant of the Fund for Scientific Research, Flanders, is grateful to the organisers for their financial support to attend the conference. The financial support of the Tournesol grant T99/040, the IUAP Contract No. P4/08 funded by the Belgian government, and the CEA, is gratefully acknowledged.

References

[1] T. Bountis, H. Segur and F. Vivaldi, Phys. Rev. A **25** 1257–1264 (1982).

[2] Chang Y. F., M. Tabor, and J. Weiss, J. Math. Phys. **23**,531–538 (1982).

[3] B. Grammaticos, B. Dorizzi and R. Padjen, Phys. Lett. A **89**, 111–113 (1982).

[4] A. P. Fordy, Physica D **52**, 204–210 (1991).

[5] S. Baker, *PhD Thesis*, University of Leeds (1995).

[6] C. Verhoeven, M. Musette, and R. Conte, J. Math. Phys. **43** (2002) 1906–1915. nlin.SI/0112030.

[7] B. Grammaticos, B. Dorizzi, and A. Ramani, J. Math. Phys. **24**, 2289–2295 (1983).

[8] J. Hietarinta, J. Math. Phys. **25**, 1833–1844 (1984).

[9] J. Hietarinta, Phys. Rep. **147**, 87–154 (1987).

[10] B. Grammaticos, B. Dorizzi, and A. Ramani, J. Math. Phys. **25**, 3470–3473 (1984).

[11] V. Ravoson, A. Ramani, and B. Grammaticos, Phys. Lett. A **191**, 91–95 (1994).

[12] F. J. Romeiras, J. Math. Phys. **36**, 3559–3565 (1995).

[13] S. V. Manakov, Zh. Eksp. Teor. Fiz. **65**, 505–516 (1973), English translation, Sov. Phys. JETP **38**, 248–253 (1974).

[14] R. Garnier, Rendiconti del Circolo Matematico di Palermo **43**, 3559–3565 (1919).

[15] S. Wojciechowski, Physica Scripta **31**, 433–438 (1985).

[16] V. G. Drinfeld and V. V. Sokolov, Proceedings of the S. L. Sobolev seminar (in Russian), **2** 5–9 (1981).

[17] S. Baker, V. Z. Enolskii, and A. P. Fordy, Phys. Lett. A **201**, 167–174 (1995).

[18] J. Satsuma and R. Hirota, J. Phys. Soc. Japan **51**, 3390–3397 (1982).

[19] A. P. Fordy and J. Gibbons, J. Math. Phys. **21**, 2508–2510 (1980).

[20] M. Błaszak and S. Rauch-Wojciechowski, J. Math. Phys. **35**, 1693–1709 (1994).

[21] V. Ravoson, L. Gavrilov, and R. Caboz, J. Math. Phys. **34**, 2385–2393 (1993).

[22] F. J. Romeiras, J. Phys. A **28**, 5633–5642 (1995).